

Yang-Baxter Equation in Spin Chains with Long Range Interactions

D. Bernard ^(a), M. Gaudin ^(a), F.D.M. Haldane ^(b) and V. Pasquier ^(a)

^(a) Service de Physique Théorique de Saclay, ¹ 91191 Gif-sur-Yvette, France.

^(b) Department of Physics, Princeton University, Princeton NJ 08544, USA.

Abstract

We consider the $su(n)$ spin chains with long range interactions and the spin generalization of the Calogero-Sutherland models. We show that their properties derive from a transfer matrix obeying the Yang-Baxter equation. We obtain the expression of the conserved quantities and we diagonalise them.

¹ Laboratoire de la Direction de Sciences de la Matière du Commissariat à l'énergie atomique.

1. Introduction.

The most remarkable properties of the XXX chain with long range interactions [1] [2] are that its spectrum is additive and that the states are created by filling a “Dirac sea” with particles obeying a “Generalised Pauli principle” [3]. Recently, it has become apparent [4] that the algebras underlying the symmetries of these models are the Yangians [5]. In [4], the first generators of the Yangian had been obtained, and the aim of this paper is to display the full algebra. To characterise it, we have constructed a transfer matrix which satisfies the exchange relations[6] resulting from the Yang-Baxter[7] equation, (often called the $RLL = LLR$ relations[8]). In the limit of infinite separation of the sites, the transfer matrix reduces to the usual XXX chain transfer matrix[7]. In proving the exchange relations, the differential operators defined in [9] [10] have turned out to be essential tools.

In the course of this work, we have been led to consider² models for which the lattice sites are replaced by dynamical particles, see also [11]. They are the spin generalisations of the Calogero-Sutherland models[12] [13]. In these models, there also exists a transfer matrix obeying the Yang-Baxter equation. However, an important difference between the two situations is that the transfer matrix of the dynamical models always commutes with the Hamiltonian, whereas in the lattice model case, it commutes only if the lattice is translation invariant. The generating function for the Hamiltonians is not given by the trace of the transfer matrix, because this trace does not commute with the Yangian. In the dynamical case it is given by the quantum determinant. In the lattice case, the determinant is a c-number which contains enough information to recover the spectrum.

In part two we define the dynamical models and we present their Lax pair. In part three we prove the Yang-Baxter equation for the transfer matrix. In part four, we derive the conserved quantities and we obtain their eigenvalues. Finally, we consider the spin models and we decompose the $su(2)$ -spin chain into irreducible Yangian representations.

2. The dynamical models.

The dynamical models are $su(p)$ generalisations of the Calogero-Sutherland model. There are M particles interacting by long range forces. Their positions are parameterised by complex numbers z_i , $i = 1, \dots, M$, and each particle carries a spin σ with p possible

² This came through discussions with M. Douglas.

values. If the particles are on the unit circle, we take $z_j = \exp(i\nu_j)$, and if they are on the line we take $z_j = \exp \nu_j$, with ν_j real. Their dynamics are governed by the Hamiltonian :

$$H_D = \sum_{j=1}^M (z_j \partial_{z_j})^2 + \sum_{i \neq j} \lambda (P_{ij} + \lambda) h_{ij} \quad (2.1)$$

where λ is a coupling constant and P_{ij} exchanges the spins of the particles i and j . The function h_{ij} is :

$$h_{ij} = \frac{z_i z_j}{(z_i - z_j)(z_j - z_i)} \quad (2.2)$$

Integrability is guaranteed by the existence of a Lax pair. It consists of two matrices L_{ij} and M_{ij} with operator entries obeying:

$$\left[H_D, L_{ij} \right] = \sum_k \left(L_{ik} M_{kj} - M_{ik} L_{kj} \right) \quad (2.3)$$

A possible choice is given by:

$$\begin{aligned} L_{ij} &= \delta_{ij} z_j \partial_{z_j} + (1 - \delta_{ij}) \lambda \theta_{ij} P_{ij} \\ M_{ij} &= -\delta_{ij} 2\lambda \sum_k (h_{ik} P_{ik}) + 2\lambda (1 - \delta_{ij}) h_{ij} P_{ij} \end{aligned} \quad (2.4)$$

with:

$$\theta_{ij} = \frac{z_i}{z_i - z_j} \quad (2.5)$$

We denote by X_j^{ab} , $a, b = 1, \dots, p$, the matrices which act as $|a\rangle\langle b|$ on the spins of the j^{th} particle and leave the other particles untouched. Using equation (2.3) and the fact that $[H_D, X_j^{ab}] = \sum_k X_k^{ab} M_{kj}$ and $\sum_j M_{ij} = 0$, one deduces that the quantities defined by:

$$T_n^{ab} = \sum_{ij} X_i^{ab} (L^n)_{ij} \quad (2.6)$$

commute with the Hamiltonian H_D . Here L^n denotes the n^{th} power of the matrix L_{ij} . Since the T_n^{ab} do not commute with each other, the spectrum is degenerate. We study their algebra in the next section.

3. The transfer matrices.

In order to arrange the algebra of the T_n^{ab} 's, we introduce the transfer matrix $T(u)$ obeying the Yang-Baxter equation:

$$R_{00'}(u-v) T^0(u) T^{0'}(v) = T^{0'}(v) T^0(u) R_{00'}(u-v) \quad (3.1)$$

According to standard notations [8], $T^0(u)$ denotes the matrix $T(u) \otimes 1$ and $T^{0'}(v)$ the matrix $1 \otimes T(v)$. The matrix $R(u)$ is the solution of the Yang-Baxter equation given by:

$$R(u) = u + \lambda P_{00'} \quad (3.2)$$

where $P_{00'}$ is the permutation operator which exchanges the two auxiliary spaces 0 and 0'. Equation (3.1) expresses the non-commutativity of the operator matrix elements of $T(u)$. The expression which we obtain for $T^0(u)$ is given by:

$$T^0(u) = 1 + \lambda \sum_{i,j=1}^M P_{0i} \left(\frac{1}{u-L} \right)_{ij} \quad (3.3)$$

where L is the matrix defined in eq. (2.4). If we set $T^0(u) = \sum_{a,b=1}^p X_{ba}^0 T^{ab}(u)$, and expand it in powers of $\frac{1}{u}$, we find:

$$T^{ab}(u) = \delta^{ab} + \sum_{n=0}^{\infty} \frac{\lambda}{u^{n+1}} T_n^{ab} \quad (3.4)$$

where the T_n^{ab} 's have been defined in eq. (2.6). We can motivate this expression as follows: It commutes with the Hamiltonian H_D , the $\frac{1}{u^2}$ coefficient coincides with (a slight generalization of) the generators of the Yangians identified in [4], and, as we will discuss in a next section, in a specific limit it gives back the transfer matrix of an XXX chain. When the z_i which define the functions θ_{ij} in equation (2.5) are complex numbers of modulus one, an important property of this transfer matrix is its hermiticity: $T^{ab\dagger}(u) = T^{ba}(\bar{u})$. As a result, the induced Yangian representation is fully decomposable into a direct sum of irreducible representations.

In order to prove (3.1), we use the formalism introduced in [9][10]. Let us define the three permutation groups: Σ_1 , Σ_2 and Σ_3 respectively generated by K_{ij} , P_{ij} and the product $(P_{ij}K_{ij})$. The operator K_{ij} exchanges the positions z_i , and P_{ij} exchanges

the spins at positions i and j : i.e. $K_{ij}z_j = z_i K_{ij}$ and $P_{ij}\sigma_j = \sigma_i P_{ij}$. We define the differentials:

$$D_i = z_i \partial_{z_i} + \lambda \sum_{i \neq j} \theta_{ij} K_{ij} \quad (3.5)$$

They obey the relations:

$$\begin{aligned} K_{ij} D_i &= D_j K_{ij} \\ [D_i, D_j] &= \lambda(D_i - D_j) K_{ij} \end{aligned} \quad (3.6)$$

We also define a projection π from the cross product $\Sigma_2 \triangleright \Sigma_3$ into Σ_2 by $\pi(\sigma\sigma') = \sigma$ for $\sigma \in \Sigma_2$ and $\sigma' \in \Sigma_3$. We use it to eliminate the permutations of Σ_1 by replacing them with those of Σ_2 . For example: $\pi(K_{12}) = \pi(P_{12}(P_{12}K_{12})) = P_{12}$. One can view this projection as the result of acting on symmetric functions under permutations in Σ_3 . The expression (3.3) of the transfer matrix then takes the following form:

$$T^0(u) = \pi \left(1 + \lambda \sum_{i=1}^M \frac{P_{0i}}{u - D_i} \right) \quad (3.7)$$

which results from $\pi(D_i^n) = \sum_j L_{ij}^n$. Since $T^0(u)$ is invariant under simultaneous permutations of the spins and the coordinates, a product of projections can be replaced by the projection of the product. Therefore, we omit the symbol π and set equal to one any permutation of Σ_3 appearing to the right of an expression. Equation (3.1) then recasts into the form :

$$\begin{aligned} & (u - v + \lambda P_{00'}) \left(\sum_{i=1}^M \left(1 + \frac{\lambda P_{0i}}{u - D_i} \right) \left(1 + \frac{\lambda P_{0'i}}{v - D_i} \right) + \sum_{i \neq j} \left(\frac{\lambda P_{0i}}{u - D_i} \right) \left(\frac{\lambda P_{0'j}}{v - D_j} \right) \right) \\ &= \left(\sum_{i=1}^M \left(1 + \frac{\lambda P_{0'i}}{v - D_i} \right) \left(1 + \frac{\lambda P_{0i}}{u - D_i} \right) + \sum_{i \neq j} \left(\frac{\lambda P_{0'j}}{v - D_j} \right) \left(\frac{\lambda P_{0i}}{u - D_i} \right) \right) (u - v + \lambda P_{00'}) \end{aligned} \quad (3.8)$$

First consider the sum over i : each term in the sum is a Yang-Baxter equation for an elementary transfer matrix, $T_i(u) = 1 + \frac{\lambda P_{0i}}{u - D_i}$, with a spectral parameter equal to $u - D_i$. The equality is therefore satisfied by each term independently. Then, consider the sum with $i \neq j$. Using the identities:

$$\begin{aligned} P_{00'} P_{0i} P_{0'j} &= K_{ij} P_{0i} P_{0'j} (K_{ij} P_{ij}) \\ P_{0i} P_{0'j} P_{00'} &= K_{ij} P_{0'i} P_{0j} (K_{ij} P_{ij}) \end{aligned} \quad (3.9)$$

the remaining terms reduce to :

$$(u - v + \lambda K_{ij}) \left(\frac{1}{v - D_i} \right) \left(\frac{1}{u - D_j} \right) = \left(\frac{1}{u - D_j} \right) \left(\frac{1}{v - D_i} \right) (u - v + \lambda K_{ij}) \quad (3.10)$$

or equivalently,

$$(u - v + \lambda K_{ij})(u - D_j)(v - D_i) = (v - D_i)(u - D_j)(u - v + \lambda K_{ij}) \quad (3.11)$$

This results from eq. (3.6). It proves the Yang-Baxter equation (3.1) for the transfer matrix given in eq. (3.3).

Finally, let us discuss the rational limit of the preceeding results. If we set $z_j = \exp(\gamma \nu_j)$ and let γ go to zero, the Hamiltonian becomes:

$$H_D = - \sum_{j=1}^M (\partial_{\nu_j})^2 + \sum_{i \neq j} \frac{\lambda(P_{ij} + \lambda)}{(\nu_i - \nu_j)^2} \quad (3.12)$$

In order to reach this limit, we need to rescale D_j and λ by γ . When γ goes to zero, the T_n^{ab} algebra degenerates to:

$$[T_n^{ab}, T_m^{cd}] = \delta_{bc} T_{n+m}^{ad} - \delta_{ad} T_{n+m}^{cb} \quad (3.13)$$

In that case, the traces $\sum_a T_n^{aa}$ define the conserved quantities which commute with the T_n^{ab} .

4. The conserved quantities.

One usually generates the conserved quantities by means of the trace of the transfer matrix. This defines quantities which commute among themselves but not with the T_n^{ab} . Here, in order to satisfy this condition, we are led to take the quantum determinant as the generating function. It is given by [6][5]:

$$Det_q T(u) = \sum_{\sigma \in \Sigma_p} \epsilon(\sigma) T_{1\sigma_1}(u - (p-1)\lambda) T_{2\sigma_2}(u - (p-2)\lambda) \cdots T_{p\sigma_p}(u) \quad (4.1)$$

The first two non-trivial terms produce the momentum and the Hamiltonian:

$$\begin{aligned} Det_q T(u) = & 1 + \frac{\lambda}{u} M + \frac{\lambda}{u^2} \left(\sum_{j=1}^M z_j \partial_{z_j} + \lambda \frac{M(M-1)}{2} \right) \\ & + \frac{\lambda}{u^3} \left(H_D + \lambda(M-1) \sum_{j=1}^M z_j \partial_{z_j} + \lambda^2 \frac{M(M-1)(M-2)}{6} \right) + \cdots \end{aligned} \quad (4.2)$$

We now give an alternative description of the transfer matrix which makes the actual computation of the quantum determinant much simpler. Let us modify the differentials D_i introduced in (3.5) as follows:

$$\begin{aligned}\widehat{D}_i &= D_i - \lambda \sum_{j < i} K_{ij} \\ &= z_i \partial_{z_i} + \lambda \sum_{j > i} \theta_{ij} K_{ij} - \lambda \sum_{j < i} \theta_{ji} K_{ij}\end{aligned}\tag{4.3}$$

Unlike the D_i , the differentials $\widehat{D}_i$ commute:

$$\left[\widehat{D}_i, \widehat{D}_j \right] = 0\tag{4.4}$$

In term of these differentials, we can define a transfer matrix which obviously satisfies the Yang-Baxter equation. It is given by:

$$\widehat{T}^0(u) = \left(1 + \frac{\lambda P_{01}}{u - \widehat{D}_1}\right) \left(1 + \frac{\lambda P_{02}}{u - \widehat{D}_2}\right) \cdots \left(1 + \frac{\lambda P_{0M}}{u - \widehat{D}_M}\right)\tag{4.5}$$

The right hand side of (4.5) is the co-product of elementary transfer matrices $\widehat{T}_i$ with their spectral parameters equal to $u - \widehat{D}_i$. It is not difficult to show that the projection of $\widehat{T}^0$ by π , $\pi(\widehat{T}^0)$, is invariant under Σ_3 . This imposes the choice of the coefficients of P_{0i} to be λ in equation (4.5). Thus, when we apply the projection π to the Yang-Baxter equation (3.1), we can replace the projection of the product $\widehat{T}^0 \widehat{T}^{0'}$ by the product of the projections. Therefore, $\pi(\widehat{T}^0)$ also satisfies the Yang-Baxter equation. Indeed, one can check that it coincides with the definition of T^0 in (3.7).

The quantum determinant of $\widehat{T}$ is the product of the quantum determinants of the $\widehat{T}_i$. It is equal to:

$$Det_q \widehat{T}(u) = \frac{\widehat{\Delta}_M(u + \lambda)}{\widehat{\Delta}_M(u)}\tag{4.6}$$

where:

$$\widehat{\Delta}_M(u) = \prod_{i=1}^M (u - \widehat{D}_i)\tag{4.7}$$

Using a similar argument to that outlined above, the invariance of $\widehat{\Delta}_M(u)$ under Σ_1 allows its replacement by its projection $\Delta_M(u)$ in (4.6) to get the quantum determinant of $T(u)$. Note that the polynomial transfer matrix $\Delta_M(u)T(u)$ also satisfies the Yang-Baxter equation.

The coefficients C_p of the polynomial $\Delta_M(u)$ define a complete set of commuting operators. We conjecture that they are given by the following ‘‘Wick’’ expansion:

$$\begin{aligned} C_p &= \sum_{i_1 < \dots < i_p} \pi(\hat{D}_{i_1} \hat{D}_{i_2} \dots \hat{D}_{i_p}) \\ &= \sum_{i_1 < \dots < i_p} \sum_{I \cup J = \{i_1, \dots, i_p\}} F_J \prod_{i \in I} z_i \partial_{z_i} \end{aligned} \quad (4.8)$$

with:

$$F_J = \sum_{\cup \{j_k, j'_k\} = J} \prod_k \left(-\lambda(\lambda + P_{j_k j'_k}) h_{j_k j'_k} \right) \quad (4.9)$$

In (4.8) and (4.9), the symbol $\cup$ stands for the union of sets which do not intersect. F_J is equal one if $J = \emptyset$ and to zero if the number of elements of J is odd. The functions h_{ij} are defined in equation (2.2). When there is no spin dependence (Calogero-Sutherland model) $\Delta_M(u)$ can be expressed as a determinant and coincides with the invariants given in [12].

Let us obtain here the eigenvalues of $\Delta(u)$. Our method is first to diagonalise the differentials $\hat{D}_i$. In this way we find the eigenvectors of $\hat{\Delta}(u)$. Then we symmetrise these eigenvectors with respect to Σ_3 to obtain the eigenvectors of $\Delta(u)$ with the same eigenvalue. In order to do this, it is convenient to make a gauge transformation which amounts to substitute $z_i \partial_{z_i} - \frac{\lambda}{2} \sum_{j \neq i} \frac{z_i + z_j}{z_i - z_j}$ for $z_i \partial_{z_i}$ everywhere. Hereafter, we indicate with a prime the gauged transformed quantities. The gauge transformed differentials $\hat{D}'_i$ take the following form:

$$\hat{D}'_i = z_i \partial_{z_i} + \lambda \left(\frac{M+1}{2} - i \right) + \lambda \sum_{j > i} \theta_{ij} (K_{ij} - 1) - \lambda \sum_{j < i} \theta_{ji} (K_{ij} - 1) \quad (4.10)$$

Their action leaves invariant the space of polynomials:

$$\Psi_{[n]}(z_1, z_2, \dots, z_M) = z_1^{n_1} z_2^{n_2} \dots z_M^{n_M} \quad (4.11)$$

where $[n]$ is a sequence of positive integers. To a sequence $[n]$, we associate the partition $|n|$ where we arrange the n_k in decreasing order. We define an order on the partitions by saying that $|n|$ is larger than $|n'|$ if the first non vanishing difference $n_k - n'_k$ is positive. It follows from the same argument as in [13] that the differentials $\hat{D}'_i$ are represented by block triangular matrices in the basis $\Psi_{[n]}(z_k)$. Namely:

$$\hat{D}'_i \Psi_{[n]}(z_k) = \sum_{n'} (d_i)_{[n'] [n]} \Psi_{[n']} (z_k) \quad (4.12)$$

with $(d_i)_{[n']|n] = 0$ if $|n'|$ is larger than $|n|$. Therefore, the eigenvalues of the differentials $\widehat{D}'_i$ are given by the eigenvalues of the block matrices on the diagonal. Let us consider such a block: $d_i^{[n]} = ((d_i)_{[n']|n'])$ with $|n|$ a fixed partition $|n| = (n_1 \geq n_2 \geq \dots \geq n_M)$. In the basis $|n_{\sigma_1}, n_{\sigma_2}, \dots, n_{\sigma_M} \rangle$, the $d_i^{[n]}$ take the following form:

$$d_i^{[n]} |n_{\sigma_1}, \dots, n_{\sigma_M} \rangle = \left(n_{\sigma_i} + \lambda \left(\frac{M+1}{2} - i + \sum_{j>i} X_{ij} - \sum_{j<i} X_{ji} \right) \right) |n_{\sigma_1}, \dots, n_{\sigma_M} \rangle \quad (4.13)$$

where the X_{ij} are defined by:

$$X_{ij} | \dots, n_{\sigma_i}, \dots, n_{\sigma_j}, \dots \rangle = \begin{cases} - | \dots, n_{\sigma_i}, \dots, n_{\sigma_j}, \dots \rangle & \text{if } n_{\sigma_i} > n_{\sigma_j} \\ 0 & \text{if } n_{\sigma_i} = n_{\sigma_j} \\ | \dots, n_{\sigma_j}, \dots, n_{\sigma_i}, \dots \rangle & \text{if } n_{\sigma_i} < n_{\sigma_j} \end{cases} \quad (4.14)$$

The matrices $d_i^{[n]}$ are triangular when we order the states inside a block by saying that $[n']$ is larger than $[n'']$ if the last non vanishing difference $n'_k - n''_k$ is positive. With the global order induced by this choice the matrices $\widehat{D}'_i$ are also triangular. It follows that the eigenvalues, $\delta_i^{[n]}$, of $d_i^{[n]}$ are the diagonal matrix elements, $(d_i)_{[n]|n]}$. One readily sees that the multiplets of eigenvalues, $(\delta_i^{[n]})_{i=1,M}$, of the $d_i^{[n]}$'s are all obtained by permuting the components of the multiplet:

$$(\delta_i^{[n]}) = \left(n_i + \lambda \left(i - \frac{M+1}{2} \right) \right) \quad (4.15)$$

As a result, the corresponding eigenvectors of $\widehat{D}'_i$ form a degenerate set of eigenvectors of $\widehat{\Delta}'(u)$ with the eigenvalue:

$$\delta^{[n]}(u) = \prod_{j=1}^M \left(u - n_j - \lambda \left(j - \frac{M+1}{2} \right) \right) \quad (4.16)$$

These eigenvectors form a representation of Σ_1 , isomorphic to the obvious representation of the permutations on the sequences $[n]$. To obtain the eigenvectors of $\Delta(u)$, one must combine these eigenvectors with a spin component and symmetrise the tensor product with respect to Σ_3 . The eigenvectors associated to a given partition $|n|$ probably organise into an irreducible representation of the Yangian.

We refer to [14] for a complementary discussion of the spectrum in the Calogero-Sutherland models.

5. The spin model.

Our understanding of the spin models[1][2] is less clear. In particular, we have not yet been able to find a simple generating function for their conserved quantities. One can find a mapping from the $su(p)$ dynamical model at $\lambda = 1$ onto the $su(p+1)$ spin chain which extends the known mapping from the Calogero-Sutherland model onto the $su(2)$ spin chain. However, the computations are cumbersome and shed no light on the transmutation of $su(p)$ into $su(p+1)$. In this section, we use the transfer matrix formalism to deduce the decomposition of the $su(2)$ spin chain into irreducible Yangian representations. From this knowledge, one can easily reconstruct the spectrum and its degeneracies.

The Hamiltonian of the $su(p)$ spin model is:

$$H_S = \sum_{i \neq j} h_{ij}(P_{ij} - 1) \quad (5.1)$$

The indices i, j refer to the sites of the chain which we take of length N . The function h_{ij} is the same as above, eq. (2.2).

The transfer matrix is the limit $\lambda \rightarrow \infty$, $\frac{u}{\lambda} = x$ fixed, of the matrix (3.3):

$$T^0(x) = 1 + \sum_{i,j=1}^N P_{0i} \left(\frac{1}{x - L} \right)_{ij} \quad (5.2)$$

with $L_{ij} = (1 - \delta_{ij})\theta_{ij}P_{ij}$. If we let the distance between the sites go to infinity with $z_1 \ll z_2 \ll \dots \ll z_N$, the θ_{ij} converge to the step function $\theta(i - j)$ and the transfer matrix reduces to its usual form:

$$T^0(x) = \left(1 + \frac{P_{01}}{x}\right) \left(1 + \frac{P_{02}}{x}\right) \dots \left(1 + \frac{P_{0N}}{x}\right) \quad (5.3)$$

For generic values of the complex numbers z_j , the induced representation of the Yangian algebra $Y(sl_p)$ is irreducible. Thus, its quantum determinant is a c-number which we can evaluate on any vector. Choosing the vector with all spins σ_j equal to p gives:

$$Det_q T(x) = 1 + \sum_{i,j=1}^N \left(\frac{1}{x - \Theta} \right)_{ij} = \frac{\Delta_N(x+1)}{\Delta_N(x)} \quad (5.4)$$

Here Θ is the $N \times N$ matrix with matrix elements θ_{ij} and $\Delta_N(x)$ is its characteristic polynomial:

$$\Delta_N(x) = \det(x - \Theta) \quad (5.5)$$

Although the transfer matrix satisfies the exchange relations (3.1) for all values of the parameters z_i , it commutes with the Hamiltonian (5.1) only if $\sum_j h_{ij}(\theta_{ij} - \theta_{ji}) = 0$ [4]. In order to satisfy this condition, we choose either $z_k = \exp(i2\pi k/N)$ with $k = 1, \dots, N$, (trigonometric models), or, $z_k = \exp(\gamma k)$, with k integers, (hyperbolic models). Diagonalising Θ , we obtain the following expression for Δ_N in the trigonometric case:

$$\Delta_N(x) = \prod_{j=1}^N \left(x - \frac{N+1}{2} + j \right) \quad (5.6)$$

We recall that in the trigonometric case, the Yangian algebra is completely reducible. Let us obtain here its decomposition in the $su(2)$ case. It is known that irreducible representations of the $su(2)$ Yangians are characterised by a polynomial $P(x)$ [5][15]. Namely, in a canonical normalisation, the transfer matrix, $T_C(x)$, is a rational function of x and one can find a state, $|\Omega\rangle$, such that:

$$T_C(x)|\Omega\rangle = \begin{pmatrix} |\Omega\rangle & 0 \\ * & \frac{P(x+1)}{P(x)}|\Omega\rangle \end{pmatrix} \quad (5.7)$$

In each irreducible block appearing in the decomposition, the transfer matrix $T(x)$ differs from the canonical one only by a multiplicative factor, $T(x) = \varphi(x)T_C(x)$. Let us evaluate the quantum determinant of $T(x)$ in this block:

$$\frac{\Delta(x+1)}{\Delta(x)} = \frac{P(x+1)}{P(x)} \varphi(x) \varphi(x-1) \quad (5.8)$$

Using the fact that $\Delta(x)T(x)$ is polynomial in x , this equation implies that the roots, $\{a_k\}_{k=1}^p$, of $P(x)$ are among those of $\Delta(x)$. If we denote by $\{b_j\}_{j=1}^{N-p}$ the remaining roots of $\Delta(x)$, it admits the solution:

$$\varphi(x) = \prod_{j=1}^{\frac{N-p}{2}} \frac{(x - b_{2j-1} + 1)}{(x - b_{2j-1})}, \quad P(x) = \prod_{k=1}^p (x - a_k) \quad (5.9)$$

only if the b_j come in disjoint pairs ($b_{2j-1}, b_{2j} = b_{2j-1} + 1$). Let us label the intervals separating consecutive roots of $\Delta(x)$ by a 1 if the extremities of the interval are b_{2j-1}, b_{2j} for some j , and by a 0 if not. Since the intervals are disjoint, two 1 can never be adjacent. If we add a 0 at both extremities of the sequence, we code the possible irreducible representations appearing in the decomposition by a sequence of $N+1$ symbols 0 and 1. In this way, we recover the characterisation of the spectrum in terms of *motifs* given in [4]. Namely, a

motif is a series of q consecutive 0 bordered by 1's; it corresponds to the natural spin $\frac{q-1}{2}$ representation $su(2)$. The representation content of a sequence is then the irreducible tensor product of its *motifs*.

Acknowledgments

We thank M.Douglas for an illuminating discussion.

References

- [1] F.D.M. Haldane, Phys.Rev.Lett. 60 (1988) 635.
- [2] B.S. Shastry, Phys.Rev.Lett. 60 (1988) 639
- [3] F.D.M. Haldane, Phys.Rev.Lett. 67 (1991) 937
- [4] F.D.M.Haldane, Z.N.C.Ha, J.C.Talstra, D.Bernard and V.Pasquier, Phys.Rev.Lett. 69 (1992) 2021.
- [5] V.G.Drinfel'd, "*Quantum Groups*", Proc. of the ICM, Berkeley (1987) 798
- [6] E.K.Sklyanin, Funct.Anal.Appl. 16 (1983) 263
- [7] R.J.Baxter, "*Exactly solved models in statistical mechanics*" Academic, London (1982)
- [8] L.D. Faddeev, "*Integrable models in 1+1 dimensional quantum field theory*" Les Houches lectures, Elsevier Science Publishers (1984)
- [9] A.P.Polychronakos, Phys.Rev.Lett. 69 (1992) 703
- [10] L.Brink, T.H.Hansson and M.Vasiliev, Phys.Letters B286 (1992) 109.
- [11] J.Minahan and A. Polychronakos, CU-TP-566 (1992) preprint ;
K.Hikami and M.Wadati, preprint (1992).
- [12] P.Calogero, J. Math. Phys. 10 (1969) 2191 ;
P. Calogero, O. Ragnisco and C. Marchioro, Lett. Nuovo Cim. 13 (1975) 383.
- [13] B.Sutherland, Phys.Rev. A5 (1972) 1372, and reference therein.
- [14] M.Gaudin, Saclay preprint SPhT/92-158.
- [15] V.Chari and A.Pressley, J.Reine Angew.Math. 417 (1991) 87.